



---

**UNDP/GEF PROJECT ENTITLED “REDUCING ENVIRONMENTAL STRESS IN THE  
YELLOW SEA LARGE MARINE ECOSYSTEM”**

---

UNDP/GEF/YS/RWG-F.4/5  
Date: 26 September 2007  
English only

**Fourth Meeting of the Regional Working Group  
for the Fisheries Component**  
*Seokcho, Republic of Korea, 7 - 9 November 2007*

**2007 Completed and On-going Activities  
Of the Fisheries Component**

1. In addition to the Symposiums on Sustainable Mariculture, Carrying Capacity and Disease diagnosis, prevention in 2007, the Fisheries Component considered implementing other activities, some of which are still pending. These activities are described below.

**Agenda 5.1 Stock Assessment**

2. The report on stock assessment has been finalised
3. The Chair (Dr Jin) will present the outcomes of the report and recommendations on how these ideas should be implemented.
4. RWG-F members are invited to review the document ([Annex 1](#)) and discuss how to implement some of the ideas presented.

**Agenda 5.2, 5.3 & 5.4 Sustainable Mariculture, Mariculture Carrying Capacity, and Mariculture Disease Diagnosis, Control and Prevention**

5. RWG-F members are invited to review the discussion document and make suggestions as to how follow up.
6. The Chair will call upon PMO to present the outcomes from the Sustainable Mariculture and Disease Meetings; Dr Fang is called upon to present the report on Carrying Capacity.
7. The discussion documents and reports from the Mariculture Symposiums/Workshops are listed in [Annex 2](#).

**Agenda 5.5 Expert Exchange**

8. The Chair will call upon Dr Yeon and himself to present ideas for the expert exchange programme

**Agenda 5.6 Regional Science Conference**

9. RWG-F members are informed of the First Regional Yellow Sea Science Conference held in Hangzhou 14 -16 August.

## Annex 1



---

### UNDP/GEF PROJECT ENTITLED “REDUCING ENVIRONMENTAL STRESS IN THE YELLOW SEA LARGE MARINE ECOSYSTEM”

---

UNDP/GEF/YS/RWG-F.4/5

Date: 28 September 2007

English only

Development of joint regional stock assessment methodology  
Comprehensive report on existing stock assessment methods and suggestions for a  
comprehensive strategy for joint-regional stock assessment

by

**Xianshi JIN**

**Third Meeting of the Regional Working Group for the Fisheries Component**

*Weihai, China, 25 - 28 October 2006*

## TABLE OF CONTENTS

1	GENERAL ISSUES .....	2
1.1	Background.....	2
1.2	Description of Activities .....	3
1.3	Expected Outputs/Results.....	4
2	METHODOLOGY .....	5
2.1	Current methods of assessing fisheries stocks: Regional & International .....	5
2.1.1	<i>Population Dynamic Models</i> .....	6
2.1.2	<i>Bottom Trawl Survey</i> .....	14
2.1.3	<i>Acoustic Survey</i> .....	15
2.1.4	<i>Tagging – Recapture</i> .....	16
2.1.5	<i>Egg and Larval Survey</i> .....	16
2.1.6	<i>Ecosystem Model</i> .....	17
2.2	Location of data & info and access to each site by the public.....	18
3	MAJOR ISSUES AND PRIORITIES .....	20
4	RECOMMENDATIONS .....	21
4.1	Collection of Fisheries and Survey Data .....	22
4.2	Stock Assessment Modelling .....	24
4.3	Jointly Scientific Survey .....	24
5	FINAL REMARKS.....	24
6	LIST OF REFERENCES .....	25

## GENERAL ISSUES

Based on the tasks listed in the Statement of Work issued by the Project Management Office (PMO) of UNDP/GEF project “Reducing Environmental Stress in the Yellow Sea Large Marine Ecosystem” (Contract Code: F-I-stockassess-JX-1201No), under the sub-objective A “Stock Assessment” of the Fisheries Component, the major task is to develop methods for joint regional stock assessment for the Yellow Sea Large Marine Ecosystem.<sup>1</sup>

### **Background**

One of the problems common to each country highlighted by a Preliminary Transboundary Analysis of the Yellow Sea Large Marine Ecosystem (Feb, 2000) is the “inadequate capacity to assess the ecosystem and perform basin-scale assessments”. The approved Implementation Plan of the UNDP/GEF Yellow Sea Project, “Reducing Environmental Stress in the Yellow Sea Large Marine Ecosystem (YSLME),” lists one of the activities of the Fisheries Component as ‘developing a common methodology for joint-regional stock assessment for the Yellow Sea’ that will ultimately enable researchers to determine the condition of stocks and manage fisheries in the future.

The major objective of the task is to develop recommendations for future stock assessments of the Yellow Sea for integration into the YSLME Strategic Action Programme (SAP) and National Yellow Sea Action Plans (NYSAPs). It is hoped that this process of joint-regional stock assessment will contribute to: the continuation of the development of regionally agreed methods for observation, monitoring and sampling of the marine environment in the Yellow Sea; an enhanced co-operative mechanism for regional monitoring and observation; an upgrading of skills in basin-wide observation and monitoring; a better scientific understanding of the basin-wide marine environment/ecosystem status; and an increased mutual understanding and trust amongst the participating institutions.

The immediate objectives of this task are to collate information on methods currently used for assessing fisheries stocks in the Yellow Sea, comparing them with methods used in other regions and for different aquatic systems, where possible filling gaps, providing

---

<sup>1</sup> **Geographic Scope:** The Yellow Sea Large Marine Ecosystem is defined in this Project Document as the body of water delineated at the south, by a line connecting the north bank of the mouth of the Chang Jiang (Yangtze River) to the south side of Cheju; at the east, by a line connecting Cheju Island to Jindo Island along the coast of the ROK; and to the north, a line connecting Dalian to Penglai (on the Shandong Peninsula).

improvements and developing a set of best-practice guidelines for stock assessment using adequate tools for assessing the condition of stocks in the Yellow Sea.

### ***Description of Activities***

- Prepare a comprehensive summary of methods used for assessing fisheries stocks in the Yellow Sea. The summary should include:
  - a. A description of the contemporary and historical techniques/methods used to assess and model stocks in this region.
  - b. A description of methods used for regular, multi-species, stock assessment in other large marine ecosystems and/or in other aquatic systems.
  - c. A description of the types of data and information collected, and used in the analysis, for all these methods.
  - d. A description of the analytical and statistical methodology utilised in each model or technique.
  - e. A description of the benefits, drawbacks and difficulties of each method.
  - f. A description of the knowledge gaps in the current method of Yellow Sea stock assessment, and the barriers to obtaining data.
  - g. Recommendations for the most practical and appropriate method for regular, multi-species, stocks assessment in the Yellow Sea with suggestions on how to fill knowledge/data/information gaps in the future.
  
- Prepare recommendations for a comprehensive guideline for annual, multi-species, joint-regional stock assessment for the Yellow Sea comprising of:
  - a. Instructions for jointly-carrying out a reiterative series of multi-species stock assessments for the Yellow Sea;
  - b. Recommendations for the acquisition of data – methodology, type, etc;
  - c. Recommendations for the analysis of data – analytical and statistical models;
  - d. Recommendations for outputs – charting and graphical display of resultant data;
  - e. Recommendations on how data should be archived or managed; and

- f. Recommendations on where final results should be published, integrated into global databases and how the public can access these data and guidelines.

The guidelines should take into consideration: i) Existing national agreements; ii) Temporal changes in climatic regimes; and, iii) The geographic parameters of the Yellow Sea Project.

- A presentation of the results-to-date at a Meeting of the Regional Working Group - Fisheries, reviewing the preliminary results with all members of the RWG-F and the Project Management Office.
- Based on the comments of the RWG-F, revise the guidelines and provide a comprehensive proposal or list of recommendations for joint-regional stock assessment (highlighting the major barriers to stock assessment, and indicating the species to be assessed) to the next Regional Working Group – Fisheries meeting, for finalisation.

### ***Expected Outputs/Results***

The final product should be a set of reports as listed below:

- i. A comprehensive report on the current techniques used to model carrying capacity both locally and internationally and comparing with other methodologies, describing data types, analytical methods, highlighting the differences, benefits, drawbacks and major barriers;
- ii. A list of data and information sources to indicate the sources of the data and information collected in item (i), location of these data and information centres, conditions of access to data and information by different users;
- iii. A recommendation for undertaking an regular, multi-species, joint-regional fisheries stock assessments in the Yellow Sea Large Marine Ecosystem, highlighting the suggested frequency of assessments, potential barriers and the species to be assessed;
- iv. Project progress and final reports as requested by the PMO.

## METHODOLOGY

Methods used to carry out assignment:

- Data and information collected during the YSLME initial data/information collection activities were referred to;
- Existing local and international data were accessed through internet searches, telephone interviews, library research, visits to/communication with national fisheries offices, research institutions, government agencies and related NGOs, compare methodologies and fill in gaps in knowledge.

### ***Current methods of assessing fisheries stocks: Regional & International***

The management objective of marine fisheries resources is the attainment of maximum sustainable yield (MSY) founded on scientific advice. This advice should be based on an evaluation of the state of the fish stock, as defined by fish abundance at a specific time, combined with the mortality and growth that control the stock development. It is often essential that advice should be given, with explicit statements of its reliability where necessary, especially when a fishery is developing rapidly. Stock assessment is presently based on two major sources of data: from fisheries and scientific surveys. Historical stock levels and the catch rate by fisheries are in most cases obtained from analyses of commercial fisheries data using Virtual Population Analysis (VPA) or alternative stock number at age based models. The present state of stocks is most commonly assessed by scientific surveys. The stock assessment by scientific surveys is required at all stages of the development of a fishery, but the need for accuracy and precision is different. In an undeveloped fishery all that is generally required is a rough measure, but the requirement of accuracy and precision increases with the development of the fishery. In a very intense or overexploited fishery, estimates with high accuracy may be necessary to provide proper management advice. The scientific survey is an important tool for assessing the present state of most commercially important stocks. The indices of abundance are used to tune a VPA or other types of catch at age models (Deriso *et al.*, 1985; Hilborn and Walters, 1992, and Patterson and Melvin, 1995). The results of fish stock assessment will provide:

- An estimate of the current stock status,
- A projection of the yield, total and spawning stock biomass and recruitment for specified scenarios of fishing mortalities, and
- The relationship between the stock status / projection and a number of biological reference points.

These parts are used to formulate biological advice on fishery management, and evaluate whether the stock is within safe biological limits, i.e. productivity (growth, recruitment) is not adversely affected by fishing.

The stock status is defined by the:

- Stock size, the number of fish by age group at a particular point in time,
- Stock productivity, growth, maturity, fecundity and recruitment, and
- Stock mortality, made up of fishing and natural mortality rates.

## **Population Dynamic Models**

To describe population dynamics of fish stocks, many models have been developed and applied world wide, the main references are the books and/or manuals by Beverton and Holt (1957), Ricker (1958, 1975) and Gulland (1969, 1983). Among the most often used models are Virtual Population Analysis (VPA), simple production model (Schaefer model) and Yield per Recruit model (Beverton and Holt model). Multi-species modelling is highly multidisciplinary in nature, including fishery science, fish biology, ecology, hydrography, mathematics, statistics, economics, operations research and computer science, the more extensive the inclusion of such factors, the more complex the models. The three previously mentioned model types will be described briefly.

### **Virtual Population Analysis (VPA)**

Although many mathematical models (including the original and corrected ones) are used to conduct stock assessment all over the world, the VPA method in fish stock assessment is widely accepted, and has become a standard assessment approach within many international communities and countries, for example, the International Council for the Exploration of the Sea (ICES), CCAMLR, CECAF,

Northern Pacific stocks, Australia, New Zealand, South Africa, Argentina, Chile, Peru, etc.

Data requirement:

Age-structured data is required by VPA. Fisheries data may provide important information about catch-at-age data from fisheries statistics and biological samples taken from these catches. The state of the stock at a specific point in time is described by:

- Stock in numbers by age group (cohort)
- Mean weight per individual in the stock by age group
- Mean weight per individual in the catch by age
- Maturity proportion by age group,  $F$  is the fishing mortality, and  $M$  is the natural mortality, total mortality  $Z=F+M$

Virtual population was originally defined by Fry (1957) as the sum of fish belonging to a given year class present in the water at any given time that are destined to be captured in the fishery. The VPA model is developed by Gulland (1965)

$$\frac{1}{N} \frac{dN}{dt} = F + M \quad (1)$$

Where:

$N$  is the number of fish in a year class

$F$  is the fishing mortality, and

$M$  is the natural mortality, which is normally assumed constant during the year. Solving equation (1) for catch gives

$$C = F \frac{N(1 - e^{-(F+M)})}{F + M} \quad (2)$$

Where:

$C$  is the catch in numbers during the year. This model also defines the demographic structure of the stock provided some additional weight at age and maturity at age are given.

The above formulas are general and not subject to any discussion on their validity for describing a fishable population of some species. The term VPA-analysis was developed using the assumption that when a year class (cohort) of fish is almost fished to extinction, an arbitrary value of  $F$  could be set to estimate the number of individuals at that age. Then, assuming a constant value of  $M$ , the numbers at any age could be calculated backwards in time by the following formula

$$C_t = F_t \frac{N_{t+1}(e^{(F_t+M)} - 1)}{F_t + M} \quad (3)$$

Where:

$t$  is the current year and  $t+1$  is the next year and solving for  $F$ . Then the following version of formula (1)

$$N_t = N_{t+1}e^{(F_t+M)} \quad (4)$$

Will give the current years population numbers and the process may be continued. The error introduced by the arbitrary choice of  $F$  at an old age would be negligible when estimating the number at age in the beginning of the life of the year class.

The problems arise when we want to calculate  $F$ .  $F$  has to be solved numerically either by reference to tables or by iteration. Either method makes the calculations somewhat laborious.

Pope (1972) proposed a method to overcome these problems by assuming that catch of each age group is taken exactly half way through each year.

$$N_2 = N_1e^{-\frac{M}{2}} \rightarrow N_1 = N_2e^{+\frac{M}{2}} \quad N_3 = N_2 - C \quad N_4 = N_3e^{-\frac{M}{2}}$$

$$\text{Therefore: } N_4 = N_1e^{-M} - C * e^{-\frac{M}{2}}$$

$$\text{or} \quad N_{t+1} = N_t e^{-M} + C_t e^{\frac{M}{2}}$$

Then:

$$N_t = N_{t+1}e^M + C_t e^{\frac{M}{2}}$$

When we take  $N_{t+1}$  as present. This means that we consider a year class from the last year it is present in the catches and work-out backwards in time. As in the Gulland's VPA,  $N_{t+1}$  has two possible forms. In the first form  $C_{t+1}$  refers to the catch in year  $t+1$  only, in this case

$$N_{t+1} = \frac{C_{t+1} * Z_{t+1}}{F_{t+1}(1 - e^{-Z_{t+1}})}$$

The second form of  $N_{t+1}$  is when  $C_{t+1}$  refers to catch in year  $t+1$  and subsequent years. This is usually the case with a completely fished year class. Then

$$N_{t+1} = \frac{C_{t+1} * Z_{t+1}}{F_{t+1}}$$

We also need a formula to calculate the  $F_t$

We can use  $N_{t+1} = N_t e^{-(F+M)}$  or

$$e^{-(F+M)} = \frac{N_{t+1}}{N_t}$$

Therefore  $-(F + M) = \ln \frac{N_{t+1}}{N_t}$  or

$$F = \ln \frac{N_t}{N_{t+1}} - M$$

If we start at age 6 we can go backwards as follows:

$$N_5 = N_6 * e^M + C_5 * e^{\frac{M}{2}}$$

$$N_4 = N_5 * e^M + C_4 * e^{\frac{M}{2}}$$

$$N_3 = N_4 * e^M + C_3 * e^{\frac{M}{2}}$$

The VPA analytical model has been expanded to include multispecies interactions (Magnusson 1995, Sparre 1991) that assumed the natural mortality mainly caused by predation in the ecosystem. Therefore, multispecies VPA (MSVPA) includes estimation of the natural mortality from predator consumption. Natural mortality is separated into two components  $M = M_p + M_o$ , where  $M_o$  is a (small) constant while  $M_p$  is calculated from the estimated stock

sizes and stomach contents. This method has only been used in a few areas due to data limitation.

### Schaefer model

Assuming an equilibrium relation between catch and population biomass, the rate of change in biomass can be written

$$\frac{dB}{dt} = G(B)$$

The Schaefer model takes this function as

$$G(B) = Bk \frac{B_{\infty} - B}{B_{\infty}} = Bk \left(1 - \frac{B}{B_{\infty}}\right)$$

Where:

$B_{\infty}$  is the maximum biomass under equilibrium and  $k$  is a constant. This equation describes the logistic growth curve.

Data requirement:

- ✓ Catch data,  $Y$
- ✓ Fishing effort,  $f$
- ✓ Catchability coefficient,  $q$

Since  $F = f.q$ , where  $F$  is instantaneous rate of fishing mortality,  $f$  is fishing effort and  $q$  is a catchability coefficient, when a fishery occurs,  $Y = FB = fqB$ , then

$$\frac{dB}{dt} = Bk \left(1 - \frac{B}{B_{\infty}}\right) - fqB$$

or

$$\frac{1}{B} \frac{dB}{dt} = k \left(1 - \frac{B}{B_{\infty}}\right) - fq$$

Under conditions of equilibrium the growth is zero and the equation becomes

$$0 = \left(1 - \frac{B_e}{B_\infty}\right) - fq \quad \text{and} \quad B_e = \frac{B_\infty}{k}(k - fq)$$

Where:  $B_e$  is the equilibrium biomass.

equilibrium yield is

The  
 $Y_e = fqB_e$

or  $Y_e = fq \frac{B_\infty}{k}(k - fq)$

Then, solving this equation, the maximum yield can be obtained.

$$Y_{e \max} = \frac{k}{4} B_\infty \quad \text{and} \quad f = \frac{k}{2q}$$

If we have an equilibrium situation, this can be fitted by plotting catch per unit effort ( $u$ ) against  $f$ :

$$u = \frac{Y}{f} = qB_e$$

The

model in non-equilibrium situations (Walter, 1975) is:  $U_{i+1} = A - Cf_i$

$$Y_{\max} = \frac{k}{4} B_\infty = \frac{k^2}{4a} \quad \text{for} \quad f = k/2q \quad \text{or} \quad Y_{\max} = \frac{A^2}{4C} \quad \text{for} \quad f = A/2C$$

Where:

$$a = \frac{k}{B_\infty}, \quad A = \frac{kq}{a} \quad \text{and} \quad C = \frac{q^2}{a}$$

## Beverton and Holt Model

Another popularly used prediction model is the “*Yield per Recruit*” model developed by Beverton and Holt (1957). The starting point of this type of model is the individual fish, compared to the Schaefer type of model which regards the total stocks as the basic unit.

The calculations are usually made in terms of the yield of a single year-class of fish throughout its life, which will be, in the steady state, the same as the yield from all year-classes in one year present in the fishery. In a stock the whole life span can be divided into periods. For each period the number alive, the number caught, the number dying of natural causes, and the number surviving to the beginning of next period can be calculated, and the yield in weight can be calculated if the number caught and mean weight of individual fish is known.

Data requirement:

This process can be calculated mathematically. The parameters of the stock concerned are given below.

$N_t$  = number of fish at age  $t$

$W_t$  = average weight of fish at age  $t$

$R$  = no. of recruits, or no. of fish alive at time  $t_r$

$M$  = instantaneous natural mortality coefficient

$F$  = instantaneous fishing mortality coefficient

We first consider a time period before fishing operates. If

$t_r$  = age at recruitment      and       $t_c$  = age at capture

then       $t_r < t < t_c$

The number ( $N_t$ ) alive at time  $t$ , can be given as  $N_t = R e^{-M(t-t_r)}$

The number ( $R'$ ) alive at first capture, can be given as  $R' = R e^{-M(t_c-t_r)}$

Therefore,

$$N_t = R' e^{-(F+M)(t-t_c)}$$

$$N_t = \text{Re}^{-M(t_c - t_r) - (F+M)(t-t_c)}$$

The yield in weight is proportional to fishing mortality (F), stock size in number ( $N_t$ ) and mean weight of an individual at age t ( $\bar{W}_t$ ).

$$\text{The yield in weight caught in a short interval } \frac{dy_t}{dt} = FN_t W_t \quad dy_t = FN_t W_t dt$$

The total weight caught throughout the life span of a cohort ( $t_c$  = age at first

$$\text{capture, } t_\lambda = \text{maximum age}) \text{ is then } Y = \int_{t_c}^{t_\lambda} FN_t W_t dt$$

$W_t$  can be expressed in the form of von Bertalanffy's growth equation

$$W_t = W_\infty (1 - e^{-k(t-t_0)})^3$$

Where  $k$  is the growth constant of the species concerned. This equation can be written as

$$W_t = 1 - 3e^{-k(t-t_0)} + 3e^{-2k(t-t_0)} - e^{-3k(t-t_0)} \quad W_t = W_\infty \sum_{n=0}^3 u_n e^{-nk(t-t_0)}$$

$$u_0 = 1, u_1 = -3, u_2 = 3, u_3 = -1$$

Therefore yield can be expressed

$$Y = \int_{t_c}^{t_\lambda} FR' W_\infty e^{-(F+M)(t-t_c)} \sum_{n=0}^3 u_n e^{-nk(t-t_0)} dt$$

writing  $t-t_0 = (t-t_c) + (t_c-t_0)$  and rearranging

$$Y = FR' W_\infty \sum_{n=0}^3 u_n \int_{t_c}^{t_\lambda} e^{-(F+M+nk)(t-t_c)} e^{-nk(t_c-t_0)} dt$$

Integration gives

$$Y = FR' W_\infty \sum_{n=0}^3 \frac{u_n}{F+M+nk} e^{-nk(t_c-t_0)} (1 - e^{-(F+M+nk)(t-t_c)(t_\lambda-t_c)})$$

If  $t_\lambda$  is sufficiently large, the last term can be neglected. The yield equation becomes

$$Y = F R e^{-M(t_c - t_r)} W_{\infty} \sum_{n=0}^3 \frac{u_n e^{-nk(t_c - t_0)}}{F + M + nk}$$

Because the recruitment is unknown and often variable, the yields etc. are normally calculated as yield etc. per recruit.

$$Y / R = F e^{-M(t_c - t_r)} W_{\infty} \sum_{n=0}^3 \frac{u_n e^{-nk(t_c - t_0)}}{F + M + nk}$$

This model assumes that the stock is in the steady state, or equilibrium condition, and that recruitment is independent of parent stock size.

### **Bottom Trawl Survey**

For demersal fish, bottom trawl surveys may provide better indices of abundance than the CPUE indices from the commercial fishery. The bottom trawl survey is a traditional and most widely used method for the assessment of demersal fish stocks. Standard surveys are all built on simple equations where an observation parameter ( $d$ ) is assumed to be directly related to the true density of fish ( $D$ ). Currently, otter trawls and beam trawls are the two major types of bottom trawls used in scientific surveys. The density of fish is estimated by swept area method.

$$d = \frac{D}{a} q$$

Where:

- $a$  is the area swept by the survey trawl, is often assumed to be the area swept by the trawl's wings or doors during a standard tow, and
- $q$  is the catchability coefficient which is affected by target fish behaviour and gear operation.

Before surveying starts, sampling design should be well-defined that will increase the precision of the survey. For example, stratified sampling is usually employed, i.e. the area is divided into several strata of more uniform fish density and the mean and the

variance for each stratum is separately estimated before combining them into a whole assessment.

Data requirement:

- ✓ fish behaviour
- ✓ gear specification
- ✓ width of wings or doors of the net

#### **Acoustic Survey**

The pelagic fish stocks are normally estimated using acoustic surveys, which usually give an absolute abundance/biomass. During the acoustic surveys, catches from bottom and pelagic trawl samplings are used to identify species in order to judge the acoustic recordings and to supply information on size composition that is needed for converting the reflected energy to actual fish densities.

The estimates of abundance obtained from the acoustic method rely on the recorded reflected energy from fish as a measure of biomass density (acoustic back scattering cross section). The basic acoustic relationship is usually formulated as.

$$s_A = \rho \cdot \sigma$$

Where:

$s_A$  is the acoustic index (the area back scattering coefficient ( $m^2/n.mile^2$ )),  $\rho$  is the area density of fish, and  $\sigma$  is the average back scattering cross section per fish (conversion factor).

For acoustic surveys,  $\sigma$  relates the acoustic signals to fish density and is correlated with the size composition of fish, which normally is established from trawling.

Instrumentation in acoustic surveys thus includes the electronic instruments as well as the gear used for sampling the recorded fish.

Data requirement:

- ✓ Target strength
- ✓ Biological characteristics

- ✓ Age, length and weight

## **Tagging – Recapture**

Conventionally, the metal tags are implanted in the abdomen, dorsal fin etc. of the fish. Recaptures are obtained from scanning catches mostly from the commercial fleet. If the mortality caused by tagging is known or assumed and the tagged fish are evenly distributed in the stock, the tagged fish can be treated as a sub-population with the same population dynamics characteristics as the stock, e.g.

$$N : T = C : R$$

This implies that the relationship between total numbers caught in the fishery (C) and number of tags found in this catch (R) reflects the relationship between the total (N) and tagged (T) populations. The development of Jolly-Seber multiple mark-recapture models (Seber 1975) enables estimations of population number (with confidence intervals) and growth rates in open populations i.e. with recruitment into, and migration out of, the population.

With the develop of techniques, active acoustic tags or new data storage tags were used, a detailed picture of an individual fish's vertical and horizontal migration can be obtained, but this method need expensive equipment and manpower.

Data requirement:

- ✓ total numbers caught in the fishery
- ✓ number of tags found in catch

## **Egg and Larval Survey**

Egg and larval surveys are another important method employed to conduct stock assessment. The total number of eggs and larvae produced per year is estimated by a survey, and divided by the number of eggs spawned per female to produce the number of fish in the spawning stock, then:

$$p = \frac{E}{FR}$$

Where:

$p$  is size of spawning stock;

$E$  is total annual egg and larva production in the survey area;

$F$  is mean annual fecundity per adult female;

$R$  is the sex ratio.

The accurate of the estimate is influenced by the mortality at egg and larval stages, and the fertilizing rate, except for the sampling problems.

Data requirement:

- ✓ total annual egg and larva production
- ✓ mean annual fecundity per adult female
- ✓ sex ratio

## **Ecosystem Model**

Due to the complex marine ecosystem, many of the single species (stock) management strategies has been a failure, ecosystem-based fisheries management have been stressed recent years. As a result, more and more ecosystem models are being developed, among them, Ecopath with Ecosim (EwE) is probably the most widely used model (Christensen, et al, 2004), and is designed for straightforward construction, parameterization and analysis of mass-balance trophic models of aquatic and terrestrial ecosystems. Focus is on using the models for fisheries management, and a suite of tools are included for this aim.

EwE has three main components:

- Ecopath is a static, mass-balanced snapshot of the system, and is used to organize historical data on trophic interactions and population sizes;
- Ecosim is a time dynamic simulation module and builds dynamic predictions by combining the data with foraging arena theory;

- Ecospace – a spatial and temporal dynamic module primarily designed for exploring impact and placement of protected areas.

The Ecopath software package can be used to

- Address ecological questions;
- Evaluate ecosystem effects of fishing;
- Explore management policy options;
- Evaluate impact and placement of marine protected areas;
- Evaluate effect of environmental changes.

The basic Ecopath equations are:

- $\text{Production} = \text{catches} + \text{predation mortality} + \text{biomass accumulation} + \text{net migration} + \text{other mortality}$
- $\text{Consumption} = \text{production} + \text{unassimilated food} + \text{respiration}$

Data requirement:

- ✓ Stomach contents
- ✓ Catches
- ✓ Biomass or density of major functional groups or species
- ✓ Migration

### ***Location of data & info and access to each site by the public***

- FAO Yearbooks of Fishery Statistics
- FAO Bulletins of Fishery Statistics
- FAO Fisheries Circulars
- FAO Handbook of fishery statistics
- FAO Fisheries Reports
- FAO Fisheries Technical Papers

- China Fisheries Yearbook
- China Ocean Yearbook
- Korean Yearbooks of Fishery Statistics
- <http://www.fao.org/fi/>
- <http://www.cnfm.gov.cn/>
- <http://www.cafs.ac.cn/>
- <http://www.ysfri.ac.cn/>
- <http://www.eastfishery.ac.cn/>
- <http://www.china-fishery.net/>
- <http://www.ifishery.com.cn/>
- <http://www.china-fisheries.org/>
- <http://www.nmdis.gov.cn/>
- <http://www.ouc.edu.cn/>
- <http://www.qdio.ac.cn/>
- <http://www.momaf.go.kr/>
- <http://www.nfrdi.re.kr/>
- <http://kodc2.nfrdi.re.kr:8001/home/eng/main/index.php>
- <http://www.pknu.ac.kr/eng/>

The data sources include: fisheries dependent data (fisheries), and independent data (scientific surveys).

The fisheries data may be extracted from fisheries statistics, mostly catch statistics by major species and total landings in whole country and total fishing effort, sometimes by fleets, the area fished is not usually specified. The large number of species caught and the many fleets operating in the Yellow Sea Large Marine Ecosystem create difficulties in getting accurate figures and sufficient fisheries data for stock assessment.

The survey data are not available to the public, and it is impossible to access to any kind of survey data on stock assessment. However, the final results from surveys can be found in publications from both countries and in international journals and books.

## MAJOR ISSUES AND PRIORITIES

Currently, three countries (China, Republic of Korea, and DPR Korea) fish in the Yellow Sea region using multi-gears operated by different types of fishing boats. Japanese fishing boats also used to operate in this area in last century with high landings. It is a fact that there is no close cooperative work on stock assessment in this region, although a lot of signs indicate that many stocks are overexploited. Management measures have been taken by individual countries unsupported by strong scientific information.

The **major problems** in stock assessment in the Yellow Sea today are:

- a. The lack of agreed methods such as those used by ICES who employs VPA tuned by survey data;
- b. The availability of data at a regional level is limited, including insufficient fisheries data and survey data;
- c. The different coverage of surveys resulting in insufficient knowledge of distribution and stock status of major species;
- d. The different methods and gear used in scientific surveys without calibration leading to different results;
- e. The different objectives of surveys leading to focus on different target species;
- f. The different target species in catch statistics;
- g. The different standards in catch statistics.

Data quality and the methods used to integrate them may result in imprecise and inconsistent stock assessments. Therefore, there is a lot of work to be done before conducting stock assessment for the whole ecosystem. To overcome these barriers, priorities are suggested as below.

First step is to establish several scientific working groups, such as

- Fisheries data WG, responsible for: collection of fisheries data; data standardization, etc.

- Survey methods WG, responsible for: comparison of survey methods used by the different countries; calibration to ensure consistent survey methods are used to estimate the biomass of stocks; development of better observation tools and survey strategies.
- Fisheries biology WG responsible for; collection of fisheries biology data; data standardization; biological characteristics of major species, e.g. growth, mortality, migration and distribution, spawning, feeding, wintering, etc.
- Stock assessment WG responsible for: selecting adequate mathematical models for the fish stocks in the Yellow Sea ecosystem; prediction of stock size of commercially important species; providing sound estimates of allowable biological catch (ABC) and total allowable catch (TAC) for fisheries management.

Second step is to establish a mechanism to exchange survey data and fisheries data.

Third step is to establish a joint survey mechanism based on analysis all the historical individual surveys performed by countries from the working groups.

## **RECOMMENDATIONS**

Stock assessment is currently based on two methods, one is mathematic models dependent on fisheries data, and the other is fisheries-independent scientific surveys. Since many population dynamics models have been developed and used in different fisheries, the suitable models for specific fisheries are the best ones based on the available data. The quality of stock assessment depends on:

- a. The quality of the assessment methods and the associated strategies;
- b. An understanding of dynamics of the ecosystem, including species interaction, relationships between environment and biology;
- c. Human activity effects;
- d. Reliable data, both fisheries and survey data;
- e. Reliability of the models chosen;
- f. Knowledge of ecosystem uncertainties.

For stock assessment in the Yellow Sea region, many traditional models have been applied to analyze the population dynamics. The models listed above, including VPA, Schaefer model and yield per recruit model are more often based on the data available in the region, and these models are highly dependent on fisheries data and their precision. All methods are useful, relevant to the Yellow Sea specific fisheries in despite of the limitation of each method is suffered to the condition of assumption, data quality, and estimation of some parameters. The accuracy is highly dependent on the knowledge of fish stocks, the precision of fisheries data collected. Attempts at building an Ecopath model have been made in recent years.

The data and information collected during the past two years for the Yellow Sea LME are more general, and can't directly fit these models without further specific fisheries data and careful processing and analyzing. However, the general trends of fisheries resources can be understood from this work.

The scientific surveys are more accurate, but costly. In the Yellow Sea region, both China and R. Korea have done some fisheries monitoring surveys, the main methods used are bottom trawl and acoustic surveys, egg and larval survey as a by-product. Tagging and recapture methods are now seldom used for stock assessment compared with past years. Those surveys data as shown in the dataset collected were insufficient for any stock assessment of the Yellow Sea LME due to the limited coverage, short period of the time series and all surveys were conducted by individual countries. However, the status of the local stocks may be estimated.

The stock assessment capacity in the Yellow Sea region needs to improve, to keep up with increasing exploitation rates in the fisheries. Although there are many scientists working with this field, capacity enhancement is required to reach equivalent standards to those employed in the northeast Pacific region and ICES regions, for example, the North Sea, the Barents, the Bering Sea, George Bank, etc. At present, due to the lack of data and capacity, the following methods for regional reiterative series of joint-regional stock assessment are as recommended below.

### ***Collection of Fisheries and Survey Data***

The data collected from fisheries and scientific surveys are used in various models to assess the state and trends in the development of stocks and to predict the catches and the development of the spawning stock in the short and medium terms.

As indicated from the models, fisheries data are important (necessary) for most models. Therefore, a collection system of accurate catch and effort and biological data should be implemented. A regularly updated fisheries database should be established that can provide a time series of stock abundance and composition.

Catch per unit effort (CPUE) is an index of stock abundance, used to support the prediction models, and may be obtained from either well-defined commercial fleets or from abundance surveys using research vessels. The CPUE values are expressed in numbers-by-age per effort unit. The effort unit can be days-at-sea, trawl hours, search time, etc. Commercial CPUE data are obtained through sampling the commercial fishery catches for biological information (for example, age composition data) and linking this information with catch and effort statistics. Abundance surveys using research vessels provide these data directly, often from bottom trawl surveys expressed as numbers caught per hour trawling. It should be noted that survey data differ from commercial CPUE data.

Biological information is the fundamental components required in the stock assessment, for example the age composition, growth, length-weight relationship, prey-predator relationships, migration, stomach contents etc.

There are large uncertainties in the ecosystem, therefore it is also necessary to collect data relating to the environment inhabited by the target fish, such as temperature, salinity, primary production, plankton, etc. Some of data is used to fit the ecosystem models.

## ***Stock Assessment Modelling***

The countries surrounding the Yellow Sea should:

- Jointly select an adequate model from the existing classical mathematical models for reiterative use in stock assessment of the fish stocks in the region after testing.
- Develop new models based on the classical and new models used in other regions.
- Agree on joint prediction of stock size of commercially important species in the region.

## ***Jointly Scientific Survey***

A regular scientific survey is suggested every year, to monitor the state of major stocks in the Yellow Sea. The sampling strategy and survey design should be well-defined. Acoustic-trawl survey is recommended, based on the availability of research vessels and equipment in the countries around the region. The frequency of cruise is completely dependent on the available budget due to the high cost of using research vessels. The coverage should be large enough to cover the whole distribution of the major stocks.

## **FINAL REMARKS**

The purpose of stock assessment aims at fisheries management for sustainable development, and resources must be managed sustainably with a long-term view. Recently, ecosystem based fisheries management (EBFM) is being discussed in many areas of the world, particularly after the Reykjavik Conference on Responsible Fisheries in the Marine Ecosystem, held in Reykjavik, Iceland, October 2001. Fisheries not only reduce the exploited stock sizes, but also affect interrelated species, including predators, prey and species in competition with the target species for food resources. It is therefore important to monitor changes in the fish community as a whole, in conjunction with the exploited stock, to ensure the ecosystem is not damaged by the fishery.

Marine fishes do not respect political boundaries and fisheries science is very internationally oriented. Therefore, countries within the same region need co-operate with each other and

work together to conduct stock assessments. When the objective of fisheries management is well defined, the method and strategy of stock assessment can be determined based on the currently available data and knowledge in relation to the Yellow Sea ecosystem.

Although there are many methods used in the world, in the Yellow Sea region, an increase stock assessment capacity is required to provide a more consistent and complete framework for sustainable management in the future. Cooperation can not flourish **without a formalized network** of institutions and scientists that maintain and share the data and knowledge. Firstly, as a result of these networks, Yellow Sea fisheries resources from different sources can be used to compare surveyed areas and assessed stocks, and all parties can directly exploit developments and improvements in assessment methodology. Secondly, effort at all levels must be encouraged to establish and to improve stock assessment in a coherent way. It is important that scientists with various areas of expertise form working groups to co-ordinate research projects.

## LIST OF REFERENCES

- Anon. 1998. Methods for fishery Resources Assessment: Status and Potentials of Marine Resource and Environment Monitoring. Report from a Working Group appointed by the Research Council of Norway. 58p.
- Beverton, R.J.H. and Holt, S. 1957. On the Dynamics of Exploited fish populations. *UK Minist. Agric. Fish. Food. Fish. Invest. (Ser. 2)* 19:533p. Facsimile reprint by Chapman & Hall.
- Cadima, E.L. 2003. Fish stock assessment manual. *FAO Fish. Tech. Pap.*, ( 393): 161p.
- Christensen, V, C.J. Walters and D. Pauly. 2004. Ecopath with Ecosim: a User's Guide. Fisheries Centre Research Reports, University of British Columbia, Vancouver. 154 p.
- Gulland, J.A., 1965. Estimation of mortality rates. Annex to Arctic fisheries working group report ICES C.M. Doc. 3 (mimeo)
- Gulland, J.A. 1969. Manual of Methods for Fish Stock Assessment - Part 1. Fish Population Analysis. *FAO Manuals in Fisheries Science ( 4)* :154p.
- Gulland, J.A. 1983. Fish stock assessment: A manual of basic methods. John Wiley, New York, 223p.
- Gunderson, D.R. 1993. Surveys of Fisheries Resources. John Wiley & Sons, Inc. New York, 248p.

- Hilborn, R. and Walters, C.J. 1992. Quantitative Fisheries Stock Assessment: Choice, Dynamics and Uncertainty. New York, Chapman and Hall, 570p.
- Johannesson, K.A. and Mitson, R.B. 1983. Fisheries Acoustics: A Practical Manual for Aquatic Biomass Estimation. *FAO Fish. Tech. Pap.*, (240):249p.
- Lassen, H. and Medley, P. 2001. Virtual population analysis. A practical manual for stock assessment. *FAO Fisheries Technical Paper*. No. 400. Rome, FAO. 129p.
- Magnusson, K., 1995. An overview of the multispecies VPA - theory and applications. *Rev. Fish Biol. Fish.* 5:195-212.
- Pope, J.G. 1972. An Investigation of the accuracy of virtual population analysis using cohort analysis. *ICNAF Res. Bull.*9:65-74.
- Ricker, W.E. 1958. Handbook of computation for biological statistical of fish population. *Bull. Fish. Res. Bd. Can.*, 119: 300p.
- Ricker, W.E. 1975. Computation and interpretation of biological statistics of fish populations. *Bull. Fish. Res. Board Can.* 191:382p.
- Saville, A. 1977. Survey methods of appraising fishery resources. *FAO Fish. Tech. Pap.*,(171): 76p.
- Seber, G.A.F. (1982) The estimation of animal abundance and related parameters. 2<sup>nd</sup> Edition, Blackburn Press, 654p.
- Sparre, P., 1991. Introduction to multispecies virtual population analysis. *ICES mar. Sci. Symp.*, 193:12-21.
- Walter, G.G. 1975. Graphical methods for estimating parameters in simple models of fisheries. *J. Fish. Res. Bd. Can.* 32:2163-2168.

#### **Websites visited or interviewed**

<http://www.fao.org/fi/default.asp>

<http://nrcwg.imr.no/Reports.html>

<http://www.imr.no/>

<http://www.nfrdi.re.kr/www06/english/index.php>

<http://www.nfrdi.re.kr/www06/english/index.php>

<http://www.pknu.ac.kr/eng/03.college/fish06.aspx>

<http://www.afsc.noaa.gov/>

<http://www.pmel.noaa.gov/sebscc/index.html>

<http://www.pices.int/>

<http://www.edc.uri.edu/lme/default.htm>

<http://www.fisheries.ubc.ca/>

## Annex 2

### Sustainable Mariculture Discussion Document

This document represents the discussions that took place following the presentations given on 19<sup>th</sup> June 2007 at the “Workshop on aquaculture in the Yellow Sea for environmental and economic sustainability and commercial profitability” at the West Sea Mariculture Research Centre in Taean.

#### Polyculture

The best combination for polyculture is animals and seaweed, but for pond culture no suitable seaweeds have been identified that can withstand the summer water temperatures.

For suspended polyculture a combination of filter feeders, seaweed and abalone is the most profitable. For seabed polyculture the combination of seaweed + abalone + sea urchin + sea cucumber + scallop has been recognized as the most profitable in China. When selecting suitable organisms for polyculture and planning species combinations and density there is a need to consider:

1. Carrying capacity
2. Economic benefit to the farmers
3. Environmental impact
4. Socio-economics

The main problems for polyculture are fouling and predator control.

#### Carrying capacity

Models have been developed to estimate the mariculture carrying capacity of coastal embayments, in China this has centred on Sungo Bay. These models give density predictions for various mixtures of species that are frequently polycultured together as discussed above. The model is still undergoing testing, however, the main problem in China is non-compliance by some farmers that increase density to increase profit when others around them have complied and decreased density to the agreed stocking levels.

As farmers were reluctant to follow the guidelines on reduction in density of cultured animals, initial demonstration sites were set up in a small area, now many farmers are following after seeing the success. Two new demonstration sites in Sungo Bay were proposed one each culture area (sea bed and suspended culture areas).

It was also suggested that the government could help by changing the way coastal areas are leased for the 15-30 years to include agreements on density and the species cultured based on the carrying capacity.

In Chile problems in the lantern culture of scallop resulted in an examination of phytoplankton density and composition and the culture density was then set according to the phytoplankton density.

The effect of these releases of hatchery produced stocks of scallops, sea urchins and abalone etc on the genetic diversity of wild stock is as yet unknown for most species. For *F. chinensis* there appears to be some genetic loss. As only a few fathers are used to produce most offspring of oysters and scallops, the possibility of genetic loss is very high amongst hatchery-raised seed, as is the likelihood of this genetic loss impacting the wild population. At present there is no regulation of genetic diversity in stock enhancement, this is not the case for disease.

In ROK regulations are being prepared at the moment, while in China there are no plans, but there is protection of certain animals and habitats

Pond polyculture is more important in ROK as no extra mariculture is allowed in the sea. However the main problem is that there are few types of seaweed than can survive the summer temperatures in a pond. Although green laver may be a suitable candidate for pond polyculture as survives it the summer in recirculation systems in ROK. In China, *Ulva* is cultured in *P. japonicus* ponds to reduce the nitrogen load. In addition amphipods are grown and are the exclusive source of shrimp food for the first 2 months. The high value *Codium* and *Gracilaria* were proposed as candidate species for pond polyculture in ROK.

### **Recirculating tank production**

Recirculation systems can be advantageous over flow through tanks systems as the reduced water exchange enables better treatment of incoming/outgoing water which can reduced the likelihood of disease entering or exiting the culture facility, limits the release of nutrients into the coastal zone and reduces the effect of red tides of cultured organisms

Land based aquaculture facilities should be recirculating to reduce the environmental impact and reduce water use. However there are bottlenecks to the development of recirculation systems and these include the high initial investment cost of equipment and the cost of energy required to drive the pumps.

These barriers to the widespread adoption of recirculation technology may be over come by following the example of the EU and USA where there are strict limits on the discharge of nutrients and taxes have to be paid per unit of nutrient discharge. These charges could well act to compensate the initial cost

At present the technology for freshwater recirculation systems in ROK mean that approximately 20% of culture area has to be sedimentation area, but this varies with different flow through rates. And if the farm has good filter facilities then the area can be reduced. In order to maintain the biological filtration the water flow has to be maintained at all times and nutrient levels changed slowly.

In ROK commercial scale freshwater systems already are in use and seawater systems are being developed. While in China some big companies are using recirculation systems.

Probiotic bacteria start-up culture produced in ROK are also used in China as inoculation for biofilters. Saltwater biofilters are less efficient in nitrogen removal than comparable FW versions, however both fish and shrimp are much less sensitive to ammonia and nitrite concentrations in SW and can survive 20ppm NH<sub>4</sub> and NO<sub>2</sub>. Some FW species can also support high N levels especially those bottom living fish.

Extruded pellets that do not dissolve so easily in the water, are found to be better than moist pellets for recirculation systems as biofilters cannot cope with large amounts particulate material. Drum filters were reported to be very useful for removal of particulate material however there are some problems in cleaning and labour.

Loss of trace elements was mentioned as a problem caused by protein fractionation> However, 2% water exchange is enough to prevent changes in alkalinity.

There is no treatment of the outflow as a result of the 2% water exchange although some developments have been made. In China, some companies have used the following

techniques to treat the outflow of land based aquaculture: outflow water→pond(culture of shrimp+filtering animals) →salt field.

### **Heterotrophic shrimp culture**

There are still some problems that need to be overcome. The high cost of construction is a major obstacle to the introduction of this culture system in ROK, although many preexisting tanks can be adapted and any material can be used.

There is also the high energy/operational cost, but some cost can be reduced by using a well insulated green house with a double layered skin with air in between> Moreover the cost of heating the water can be reduced using underground water or cooling water from power plants or efficient heat exchangers. Oxygen may not be needed and could be substituted by an efficient aeration system. There is also a lack of knowledge and experience to train the farmers, through this may be supported by an implementation research project. Also in ROK there is a lack of infrastructure to provide SPF seeds or HHS larvae- good quality.

This method should be implemented in the ROK for indoor culture, as the whole can be isolated from disease. Most culture here is either successful or faces total collapse from disease problems

In China, the indoor culture of shrimp is just in small commercial scale and it has been developed slowly because of the higher cost and low price for both *F. vannamei* and *P. chinensis* (USD 3/ kg) and therefore could not make a profit. The government supplies Specific Pathogen free (SPF) juveniles, and support companies to keep broodstock indoors in biosecure systems. There are now 4 shrimp breeding government facilities that supply SPF seeds at the same price as regular seeds. SPF seed are not normally more resistant to disease than regular shrimp. Once they leave the secure facility they are now High Health Shrimp (HHS). Specific Pathogen resistance (SPR) is achieved through genetic selection.

### **Open Sea Aquaculture Technology and practices**

Until 2009 only one license per province in ROK for offshore culture demonstration sites has been issued. However one of the problems that remain is how to define offshore?

Around the world it depends on the depth and distance from shore, in ROK there is no definition yet. In China applications for offshore culture have to be submitted, and so far the definition of offshore has not been set, but >10km is being considered. At present in China longline culture and deep sea cages have to be in >15m-20m and after 5 yrs the government will encourage cage development in 20 – 40m. Also in areas near Chinese cities no aquaculture is allowed within 1500m from shore. While in ROK all coastal waters < 15m in depth belongs to the village people cooperatives, therefore all other activities have to be in deeper water. Offshore/deep water culture is not really appropriate in the Yellow Sea as Chinese farmers try to save money by placing the cages nearshore as the Yellow Sea is very shallow which would mean that deep sea cages are unpractical as they would have to be so far offshore.

The government in ROK wants to encourage the movement of nearshore cages to offshore locations by 2010 and there are also plans to move the Chinese cage culture offshore to deeper water.

Offshore cage culture is more sustainable and responsible than nearshore culture as initial results suggest:

- It is environmentally friendly as the deeper water ensures greater dispersion of any nutrient inputs and less seafloor deposition of uneaten food
- HACCP principle: food safety of the produce is improved as it is less likely to be affected by anthropogenic pollution

- Utilization of the ocean in a 3 dimensional manner
- That better quality fish produced from offshore culture commands higher prices
- There is reduced coastal pollution caused by accumulation of feedstuffs
- Improved tourism potential as visual and water quality impact of nearshore cage culture are reduced
- Better growth and survival of culture animals as a result of improved water quality

### **Best Management Practices (BMP) on finfish/shellfish culture**

In order to grow fish faster and healthier and in a more sustainable manner BMPs need to be followed. Details of BMPs were outlined in the presentation. Unstressed fish were reported to have better growth and survival.

In China fish cages are often grouped which can create environmental problems from localized enrichment but in other countries the cages culture are separated by much distance eliminating this problem.

In ROK FW licenses were issued for 10 yr periods, but due to rising nutrient levels in the culture areas no more permissions have been issued which had caused the cessation of FW cage culture. However this was probably a mistake as no improvement in water quality has been subsequently noted, suggesting that nutrients are coming from elsewhere.

For shellfish one of the major concerns in China is the Manilla clam summer mortality, caused by an influx of the freshwater from storms together with slow water movement. In order to avoid these stresses, they now practice rotation of culture sites and have expanded the culture area to the sub-tidal zone (5-6m below sea level) where the growth rates are much higher. Manilla clam was also reported to grow well in ponds.

Also of concern is the fact that clams are now cultured in areas close to cities where there is an increase in concentration of biotoxins and land-based pollution. This may make them unsafe for human consumption.

### **Best management for feed and feeding**

One of the major advances is the formulation of better diets using suitable ratios of nutrients, energy, fatty acids, amino acids and protein ratios that are adapted for each species. These features combined with improved water stability and better manufacturing techniques have resulted in higher quality diets that pollute less as the proportion of the diets energy that is excreted is less because of improved palatability water stability and digestibility. Regulation of the feeding frequency, amount and time of feeding through automatic or manual feeding can also help reduce wastage and environmental impact. The use of alternative protein and fatty acids sources can help reduce the amount of fish biomass (trash fish) used in the diet as can careful selection of the species used for aquaculture

The use of trash fish has become a major issue as not only are trash fish more polluting when used as a feed in many countries the same fish are also used for human consumption. In ROK fishermen are catching undersized fish illegally. The trash fish are normally caught offshore in coastal areas, in this region anchovy is the preferred trash fish. Compound feeds used for cage culture in China contain 30% fishmeal by dry weight, in ROK this rises to 50% dry basis for cage culture pellets. This is still much less than 100% for trash fish. Therefore it is necessary to persuade the farmers to use dry pellet but this is very difficult as trash fish appears much cheaper. In China there is a subsidy for dried pellets is 20% to encourage use and the government plans to ban the use of trash fish in 10 years in China. At demonstration sites farmers were given compound foods (pellets) and asked to compare the resulting growth and survival with fish fed trash fish, and even though the results for pellets were superior still some farmers were reluctant to use the compound food. Price is still the major issue.

In many shrimp diets the use soybean meal and poultry by product has been a research focus to help decrease the amount of fishmeal used and reduced the price of feeds. The selection of culture animal used can also have a major impact on the amount of protein (=fishmeal) that is required in the diet. For example *Penaeus vanamei* requires ~ 20% protein, *Fenneropenaeus chinensis* ~ 40% and *Penaeus japonica* > 50%.

### **Information exchange between scientists, policy makers and farmers.**

#### China

Mariculture is regulated in Central Government through the Agriculture Ministry. This ministry in combination with the Provincial Govt Ocean Fisheries Bureau operates a Fisheries Technical Extension Service to run training courses, workshops and demonstration sites. Moreover at every level from Provincial Ocean Fisheries Bureau down to City and County levels there are also mariculture and inland fisheries institutes that regulate and provide information and advice.

And institutes like the YSFRI, under the control of the Agricultural Ministry, are very closely linked to the commercial enterprises so the flow of information is very good. The government actively encourages companies and scientists to work closely.

Agricultural ministry gets advice from technical reports, project outcomes and experts.

#### R. Korea

Ministry of Mariculture and Fisheries (MOMAF) operates >10 local maritime fisheries offices with extension workers that offer small scale workshops for farmers. Bigger issues are addressed in bigger workshops

In addition the National Fisheries Research and Development Institute (NFRDI) hosts workshops every year on each species. They also operate service agents, offer information and advice through there website, produce technical guides for various species that are distributed amongst farmers, and operate demonstration sites where new technologies can be explained. There are also 7 different websites that have aquaculture/mariculture chat room where farmers and experts can exchange views

Mariculture use a self regulation systems, and every year symposiums are held for the Fisheries Cooperative (80% are fish farmers) to disseminate information which the farmer are encouraged to follow as they receive benefits from the government and have to attend the courses.

NFRDI provide advice to MOMAF and the Govt give projects to KORDI/KMI to evaluate management decisions. Korean aquaculture and fisheries societies also provide information